

A Proposal for Low Energy Dilaton

HoSeong La

*Department of Physics, North Dakota State University
Fargo, ND 58105-5566, USA*

E-mail: hsla@golem.phys.ndsu.nodak.edu

and

*Center for Theoretical Physics, Laboratory for Nuclear Science,
Massachusetts Institute of Technology, Cambridge, MA 02139-4307, USA*

We propose a systematic way of introducing the (nongravitational) low energy dilaton and a scheme for spontaneous breaking of scale symmetry (SBSS) is explained.

The dilaton is a hypothetical particle which is a Goldstone boson associated with SBSS. The earlier idea of low energy dilaton was abandoned because, in renormalizable QFT's, the scale symmetry suffers from the trace anomaly. Motivated by recent interests in the gravitational dilaton, we decided to revisit the dilaton in the nongravity context. If we look into the matter more carefully, the existence of anomalies does not necessarily rule out the possibility of SSB. If SSB occurs in a sector different from the anomalous one, it is possible. A good example is the axion. Despite the axial anomaly, spontaneous breaking of the Peccei-Quinn symmetry occurs so that the axion can be generated in QCD. Furthermore, the dilaton does not transform like a quasiprimary field under dilatations, the RG argument does not rule out the possibility of SBSS.

There are two key facts in our proposal. First, the dilaton is systematically incorporated into the low energy scale invariant effective theories by introducing the “dilaton geometry” such that the diffeomorphism (Diff) invariance in the dilaton geometry is equivalent to the conformal invariance of the Minkowski space in the presence of the dilaton. Second, we also need to generalize the scale symmetry by introducing an auxiliary scale dependent constant, hence making the scale symmetry “off shell”. This is necessary because the usual scale symmetry does not allow a potential other than the one with the runaway dilaton vacuum. Such a constant is in some sense nothing new. For example, see the Wilson's approach to RG. This talk is based on [1], in which more details can be found.

The dilaton geometry is defined by a metric of the form

$$g_{\mu\nu} = e^{2\kappa\phi} \eta_{\mu\nu}, \quad (1)$$

where $e^{4\kappa\phi} = \sqrt{|\det g_{\mu\nu}|}$ and κ^{-1} is the dilaton scale. As pointed out in [2], if $\delta\phi$ under Diff is not as a scalar, but dictated by the metric, then

$$\delta e^{\kappa\phi} = \frac{1}{2} e^{\kappa\phi} (\partial_\mu + 4\kappa \partial_\mu \phi) v^\mu. \quad (2)$$

Under volume-preserving diffeomorphisms (SDiff), ϕ behaves like a constant. It is important to notice that the consistency condition between eq.(2) and eq.(1) is

$$\frac{1}{2}\eta_{\mu\nu}\partial_\alpha v^\alpha = \eta_{\mu\alpha}\partial_\nu v^\alpha + \eta_{\alpha\nu}\partial_\mu v^\alpha. \quad (3)$$

Thus, diffeomorphisms of eq.(1) appear as conformal transformations of the flat spacetime. This shows that the dilaton geometry describes the conformal symmetry of the flat spacetime in the presence of the dilaton. Under dilatations $v^\mu = \alpha x^\mu$, the dilaton transforms inhomogeneously as

$$\delta\phi = \alpha \left(\frac{1}{\kappa} + x^\mu \partial_\mu \phi \right), \quad (4)$$

hence the dilaton is not a quasi-primary field. This distinguishes the dilaton from other fields. Nevertheless, the dilaton is a Lorentz scalar.

Let us define a dilaton-dressed field as $\Phi_{[d]} \equiv e^{d\kappa\phi}\Phi$, where Φ a scalar in the dilaton geometry. Then, under dilatations, $\delta\Phi_{[d]} = \alpha(d + x^\mu \partial_\mu)\Phi_{[d]}$ so that $\Phi_{[d]}$ is a quasiprimary field. This recovers the usual dilatations in the flat spce: $\Phi_{[d]}(x) \rightarrow e^{d\alpha}\Phi_{[d]}(e^\alpha x)$ for $x \rightarrow e^\alpha x$, where d is the scale dimension (or the conformal weight). Such dressing is not needed for vector fields in four dimensions because under dilatations $\delta A_\mu = \alpha(1 + x^\lambda \partial_\lambda)A_\mu$. This in particular leads to the YM term that does not couple directly to the dilaton in four dimensions, hence different from the Kaluza-Klein approach of introducing the dilaton. In other than four dimensions vector fields still need dilaton dressing, leading to direct YM-dilaton couplings. The mass dimension of a field is not necessarily the same as its scale dimension. For example, the dilaton has mass dimension 1, but its scale dimension is not even defined.

To define the (e.g. $\mathcal{N} = 1$) supersymmetric dilaton geometry we need a supersymmetric generalization of SDiff. As SDiff leaves the volume density $\sqrt{g}$ invariant, supersymmetric SDiff should also leave the chiral density $\mathcal{E}$ invariant. In curved spacetime, therefore, we demand $\delta(\tilde{\sigma}_{\alpha\dot{\alpha}}^\mu \psi_\mu^\alpha) = 0$. Using $\tilde{\sigma}_{\alpha\dot{\alpha}}^\mu \psi_\mu^\alpha = E_a^\mu \tilde{\sigma}_{\alpha\dot{\alpha}}^a \psi_\mu^\alpha$, we obtain $\delta(\tilde{\sigma}_{\alpha\dot{\alpha}}^\mu \psi_\mu^\alpha) = v^\mu \partial_\mu (\tilde{\sigma}_{\alpha\dot{\alpha}}^\mu \psi_\mu^\alpha) + 2\tilde{\sigma}_{\alpha\dot{\alpha}}^\mu \partial_\mu \zeta^\alpha = 0$. This defines the fermionic part of the supersymmetric SDiff. The supersymmetric dilaton geometry needs fermionic analog of eq.(1), and a good candidate is

$$\psi_\mu^\alpha = \bar{\sigma}_\mu^{\dot{\alpha}\alpha} \bar{\psi}_{\dot{\alpha}}, \quad (5)$$

where σ -matrices are those in flat space. Then these supersymmetric diffeomorphisms correctly reproduce the superconformal transformations of flat superspace. Thus, defining the supersymmetric dilaton geometry, we identify ψ as the superpartner of the dilaton, the dilatino. Its transformation rule under supersymmetric dilatations is

$$\delta\bar{\psi}_{\dot{\alpha}} = v^\mu \partial_\mu \bar{\psi}_{\dot{\alpha}} + \frac{1}{4}\bar{\psi}_{\dot{\alpha}} \partial_\mu v^\mu - \frac{1}{2}\sigma_{\alpha\dot{\alpha}}^\mu \partial_\mu \zeta^\alpha, \quad (6)$$

where $2\bar{\sigma}_\mu^{\dot{\alpha}\alpha}\partial^\mu\bar{\eta}_{\dot{\alpha}} = 2\partial^\mu\partial_\mu\zeta^\alpha = \frac{1}{2}\partial^\mu\partial_\mu\zeta^\alpha = 0$. Thus the supersymmetric dilaton geometry is an effective way of describing the supersymmetric dilatations in the flat space.

From the supersymmetric point of view, the dilaton is always associated with the axion. In the dilaton geometry the axion can be incorporated by defining $\chi = e^{\kappa\phi_c}$ such that $g_{\mu\nu} = \chi^*\chi\eta_{\mu\nu}$, where $\kappa\phi_c \equiv \kappa\phi + i\frac{a}{f_a}$. χ is mass-dimensionless. Although χ is not a scalar in the dilaton geometry, we can demand that χ should still transform like a scale-dimension one field under dilatations so that the axion transformation rule can be obtained as $\delta a = \alpha x^\mu\partial_\mu a$. This axion can be related to the antisymmetric field $B_{\mu\nu}$ in the dilaton geometry. Thus, the dilaton, axion and dilatino form a chiral supermultiplet (ϕ_c, ψ) . The R-charge of the dilaton-axion multiplet can be easily determined from this and the superconformal symmetry such that its R-charge is zero. This enables us to define dressing for fermions. Fermions should be dressed over each Weyl components such that $\Psi_{[d]} \equiv \begin{pmatrix} \psi_{1[d]\alpha} \\ \bar{\psi}_{2[d]}^{\dot{\alpha}} \end{pmatrix} = \begin{pmatrix} \chi^d & 0 \\ 0 & \chi^{*d} \end{pmatrix} \begin{pmatrix} \psi_{1\alpha} \\ \bar{\psi}_2^{\dot{\alpha}} \end{pmatrix} \equiv \begin{pmatrix} \chi^d & 0 \\ 0 & \chi^{*d} \end{pmatrix} \Psi$.

We demand a superconformally invariant action in the presence of the dilaton-axion multiplet to be covariant in the supersymmetric dilaton geometry. Thus, such an effective action can be read off from the supergravity action using the metric of the supersymmetric dilaton geometry. The bosonic case can be analogously defined too. Introducing a dynamical scale which transforms under dilatations as $\delta M = \alpha M$, hence making the scale transformation “off-shell”, we can in fact write down a dilaton effective potential which is free from a trace anomaly: A generic form of the pure dilaton part of a scalar potential which exhibits SBSS is

$$V_{\text{dil,eff}} = \kappa^2 \Lambda e^{4\kappa\phi} (\phi - a_1)(\phi - a_2), \quad (7)$$

where Λ is a mass dimension four constant and $\kappa a_i = \frac{1}{2} \ln \frac{\kappa^2 \Lambda}{M^2} + c_i$, $i = 1, 2$, and c_i 's are constants independent of M . The specific forms of a_i are dictated to make the potential scale invariant so that there is no trace anomaly from this potential. This scalar potential can be derived from a superpotential, for example, with one scalar field, z , other than the dilaton,

$$W_{\text{eff}}(\phi_c, z) = \frac{1}{2\sqrt{3}}\sqrt{\Lambda}(a_1 - a_2)e^{3\kappa\phi_c} + \sqrt{\Lambda}\kappa z e^{2\kappa\phi_c} \left(\phi_c - \frac{1}{2}(a_1 + a_2)\right). \quad (8)$$

The first term is an R-symmetry breaking term that vanishes as $(a_1 - a_2) \rightarrow 0$. $V_{\text{dil,eff}}$ has a scale symmetry breaking vacuum at

$$\langle\phi\rangle = v \equiv \frac{1}{2} \left(a_1 + a_2 - \frac{1}{2\kappa} + \sqrt{(a_1 - a_2)^2 + \frac{1}{4\kappa^2}} \right) \quad (9)$$

with $V_{\text{dil,eff}}(\langle\phi\rangle) \leq 0$. The equality is only for $a_1 = a_2$. The dilaton mass in the nonsymmetric vacuum reads $m_\phi^2 = 2\kappa^2 \Lambda e^{4\kappa v} (1 + 2\kappa(2v - a_1 - a_2))$. As

$v \rightarrow -\infty$, $m_\phi \rightarrow 0$, confirming the dilaton is massless in the scale invariant asymptotic vacuum. To be more precise, the vacuum must be a vacuum for the total scalar potential V_{eff} , not just the pure dilaton part V_{dil} . The vacuum structure of V_{eff} is fairly complicated even with just one chiral multiplet. There are three different types of vacua with SBSS: unbroken supersymmetry, spontaneously broken supersymmetry and softly broken supersymmetry. If the scale symmetry is broken in the bosonic vacuum, then the Poincaré supersymmetry must be broken at the same time. If the scale symmetry is broken in the fermionic vacuum but the bosonic vacuum remains invariant, then the Poincaré supersymmetry can be preserved as long as the R-symmetry breaking is specifically related to the scale symmetry breaking

If $a_1 = a_2$, there is no axion contribution to the scalar potential other than the axion mass term in my examples. Otherwise, however, the scalar potential explicitly contains nontrivial axion potential in some cases. The axion effective potential roughly takes the form of

$$V_{\text{eff}}(a) = f(\phi, z, \bar{z})a^2 + g(\phi, z, \bar{z})(a_1 - a_2) \\ \times \left[\text{Re}z \left(h(\phi) \cos \frac{a}{f_a} + \frac{a}{f_a} \sin \frac{a}{f_a} \right) - \text{Im}z \left(\frac{a}{f_a} \cos \frac{a}{f_a} - h(\phi) \sin \frac{a}{f_a} \right) \right]$$

for smooth functions f , g and h . In particular, the potential is not periodic. The first term is due to the term necessary for scale symmetry breaking, hence the dilaton contribution breaks the periodicity of the axion potential. The nonperiodicity in the second term is because the corresponding term in the superpotential does not have R-symmetry. Nevertheless, it still yields an extremum at $a = 0$. The periodic axion potential can be obtained in the limit $h(\phi) \sim \phi \rightarrow -\infty$ for $\phi \gg a$, which is the scale invariant limit, and z gets a real vacuum expectation value. All the vacua I obtain determine $\langle a \rangle = 0$.

If we incorporate such a structure into the gravitational case such that the low energy dilaton is related the gravitational dilaton, the Diff symmetry of curved spacetime may appear as spontaneously broken down to SDiff symmetry. Although this still contains the Poincaré symmetry, hence, nothing against current experimental observations, there is much prejudice against such an idea. We hope nature herself will clarify this in future.

Acknowledgements: I thank R. Arnowitt, D.Z. Freedman, M. Grisaru and G. Mack for helpful correspondence. I am also much grateful to B. Zwiebach at MIT and R. Hammond at NDSU for their hospitality while this work was done. This work is supported in part by DOE and NASA.

References

1. H.S. La, hep-th/9608148, hep-th/9704196.
2. H.S. La, hep-th/9510147.